


## PHYS 210 - GENERAL PHYSICS I KNOW SHEET

There are some basic tools that a scientist should have at his/her disposal without having to consult a reference. This list consists of relationships that will serve you well to know. Commit these all to memory! Items 1 - 8 you should have seen in high school mathematics at some point. Items 9 and 10, the Constants, and the Formulae we will encounter this semester - make sure you commit them to memory after they are introduced in a reading assignment.

1.  $a x^2 + b x + c = 0 \quad \Leftrightarrow \quad x = \frac{1}{2 a} [ - b \pm \sqrt{b^2 - 4 a c} ]$


2. For right triangles:  $\sin A = y/r$   
 $\cos A = x/r$   
 $\tan A = y/x$   
 $x^2 + y^2 = r^2$



3.  $\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$   
 $\cos (A \pm B) = \cos A \cos B \mp \sin A \sin B$

4. Law of Cosines:  $a^2 = b^2 + c^2 - 2 b c \cos A$

Law of Sines:  $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$



5. Circle:  $C = \text{circumference} = 2 \pi r$

$A_{\odot} = \text{Area} = \pi r^2$

6. Cylinder:  $A = \text{surface area} = 2 \pi r L + 2 \pi r^2$

$V = \text{volume} = \pi r^2 L$

7. Sphere:  $A = \text{surface area} = 4 \pi r^2$

$V = \text{volume} = 4/3 \pi r^3$

8.  $\sin^2 A + \cos^2 A = 1$

9.  $|\vec{a} \times \vec{b}| = a b \sin \theta$ ;  $\vec{a} \times \vec{b} = \hat{i} (a_y b_z - b_y a_z) + \hat{j} (a_z b_x - b_z a_x) + \hat{k} (a_x b_y - b_x a_y)$

10.  $\vec{a} \cdot \vec{b} = a_x b_x + a_y b_y + a_z b_z = a b \cos \theta$

Physical Constants: [Memorize as we encounter these in class.]

$g = 9.8 \text{ m/s}^2 \quad G = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2 \quad c = 3.00 \times 10^8 \text{ m/s}$

Physical Formulae: [Memorize as we encounter these in class.]

$\vec{v} = \frac{d\vec{r}}{dt} \quad \vec{a} = \frac{d\vec{v}}{dt} \quad \vec{F}_{NET} = m\vec{a} = \frac{d\vec{p}}{dt} \quad W = \int \vec{F} \cdot d\vec{r} \quad U_g = mgy \quad a_c = \frac{v^2}{r}$

$F_G = mg \quad \vec{F}_{GRAV} = \frac{GmM}{r^2} \hat{r} \quad F_s \leq \mu_s N \quad F_{spring} = -kx \quad K = \frac{1}{2} m v^2$

$\vec{p} = m\vec{v} \quad \vec{\tau} = \vec{r} \times \vec{F} \quad \vec{v} = \vec{\omega} \times \vec{r} \quad \vec{L} = \vec{r} \times \vec{p} \quad f = \frac{1}{T} \quad \omega = \frac{2\pi}{T}$

Text Appendix A: Derivatives: Know them all! Integrals: Know the first five in the first column and the 3<sup>rd</sup>, 5<sup>th</sup>, and 6<sup>th</sup> in the second column [integrals of  $e^{ax}$ ,  $\sin(ax)$ , and  $\cos(ax)$ ]